

# STUDIES & FORMULAE OF PROPERTIES USING ULTRASONIC VELOCITY

- AZEOTROPISM
- INTER MOLECULAR INTERACTION
- MISCIBILITY AND COMPATIBILITY
- PHASE TRANSITION
- THERMODYNAMIC PROPERTIES
- TRANSPORT PROPERTIES ETC.

## 1. Adiabatic Compressibility( $\beta_{ad}$ )

$$\beta_{ad} = (\rho v^2)^{-1}$$

where  $\rho$  = density

Ref: A. Varada Rajulu and P. Mabu Sab, Bull. Mater. Sci., Vol. 18 (June 1995), No. 3, pp. 247-253.

## 2. Isentropic Compressibility( $\beta$ )

$$\beta = (\rho v^2)^{-1}$$

where  $\rho$  = density

Ref: D.N. Rao, A. Krishnaiah and P.R. Naidu, Acta Chim. Acad. Sci. Hung, Vol. 107 (1981), No. 1, pp. 49-55.

## 3. Isothermal Compressibility( $\beta_T$ )

$$1. \quad \beta_{T,n-mix} = \frac{17.1 \times 10^{-4}}{v_{n-mix}^2 T^{4/9} \rho_{n-mix}^{4/3}}$$

where  $\rho$  = density.

Ref: Vimla Vyas, Physics and Chemistry of Liquids, Vol.42, No.3, June 2004, pp.-229-236.

$$2. \quad \beta_T = \beta_S + \frac{TV\alpha^2}{C_p}$$

where  $\alpha$  = Thermal Expansivity,  $C_p$  = heat capacity at constant pressure.

Ref: K.S.Reddy, M.Sreenivasulu and P.R.Naidu, Zeitschrift fur Physikalische Chemie Neue Folge, Bd. 124, S. 149-154 (1981).

## 4. Effective Debye Temperature ( $Q_D$ )

$$Q_D = \frac{h}{k} \left[ \frac{(9N/4\pi V)}{(P\beta_{ad})^{3/2} \left\{ \left( \frac{1}{1+\gamma} \right)^{3/2} + 2 \left( \frac{4}{3\gamma} \right)^{3/2} \right\}} \right]$$

where  $h$ = Planck's constant,  $k$ = Boltzmann's constant,  $N$ =Avogadro number,  $V$ = molar volume,  $\rho$  = density,  $\beta_{ad}$ = adiabatic compressibility and  $\gamma = \frac{\beta_T}{\beta_{ad}}$  = specific heat ratio.

Ref: *J.D. Pandey, V. Sanguri, R.K. Mishra and A.K.Singh*, J.Pure Application Ultrason, Vol. 26, 2004, pp. 18-29.

### 5. Gruneisan Parameters ( $\Gamma$ )

$$\Gamma = v^2 \alpha / C_p$$

where  $\alpha$  = thermal expansion coefficient,  $C_p$  = Principal Heat Capacity at constant pressure.

Ref: *B.P. Shukla, L.K. Jha, A.P. Upadhyay, S.N. Dubey and S.N.D. Dubey*, J. Pure Appl. Ultrason., Vol. 11 (1989), pp. 32-33.

### 6. Intermolecular Free Path Length ( $L_f$ )

$$L_f = K \sqrt{\beta_{ad}}$$

where  $\beta_{ad}$ = adiabatic compressibility,  $K$  = temperature dependent Jacobson's Constant

Ref: *P.S. Nikam and Mehdi Hasan*, Asian Journal of Chemistry, Vol. 5 (1993), No. 2, pp. 319-321.

### 7. Internal Pressure ( $\pi_i$ )

$$\pi_i = brt \left[ \frac{K\eta}{v} \right]^2 \frac{\rho^{2/3}}{M^{7/6}}$$

where  $b$ =(packing factor),  $R$ =universal gas constant,  $K$ =temperature independent constant,  $\eta$ =viscosity of liquid,  $\rho$  = density and  $M$ = molecular weight.

Ref: *C.V. Suryanarayana and P. Pugazhendhi*, Indian Journal of Pure & Applied Physics, Vol. 24 (Aug. 1986), pp. 406-407.

### 8. Free Volume ( $V_f$ )

$$V_f = \left( \frac{Mv}{K\eta} \right)^{3/2}$$

where  $K$ =temperature independent constant,  $M$ = molecular weight,  $\eta$ =viscosity of liquid,

Ref: *N. Prasad and H. Rajendra*, J. Pure Appl. Ultrason., Vol. 25 (2003), pp. 25-30.

### 9. Rao's Constant ( $R$ )

$$R = V (v)^{1/3} \quad \text{or} \quad R = (M / \rho) v^{1/3}$$

where  $\rho$  = density,  $V$  = molar volume and  $M$ = Molecular Weight.

Ref: *R. Paladhi and R.P. Singh*, Acustica, Vol. 72, (1990), pp. 90-95.

### 10. Surface Tension ( $S$ )

$$v = (S / 6.3 \times 10^{-4} \rho)^{2/3}$$

where  $\rho$  = density.

Ref: *V.K. Syal, R.Gautam and S. Chauhan*, Indian Journal of Pure & Applied Physics, Vol. 36, (1998),

### 11. Mean Square Thermodynamic Fluctuation

$$\text{Mean square fluctuations of pressure, } \overline{(\Delta P)^2} = (kT\rho v^2)/V \quad (1)$$

$$\text{Mean square fluctuations of Temperature, } \overline{(\Delta T)^2} = \frac{kT^2\gamma}{C_p} \quad (2)$$

$$\text{Number of molecular in a given volume, } \left(\frac{\Delta N}{N}\right)^2 = kT\gamma/\rho v^2 V \quad (3)$$

$$\text{Correlation function of pressure and temp. fluctuations, } \overline{\Delta P.\Delta T} = \frac{\alpha\rho v^2 kT^2}{C_p} \quad (4)$$

where k = Boltzman's constant, T = temperature in K,  $\rho$  = density,  $\gamma$  = ratio of specific heats,  $C_p$  = specific heat at constant pressure and  $\alpha$  = coefficient of expansion.

(1)-(4). Ref: *R. Sabesan, R. Varadarajan and S. Srinivasan*, Jour. Acoust. Soc. Ind., Vol. XII (July 1984), No. 3, pp. 38-40.

### 12. Van der Waal's Constant (b)

$$b = \frac{M}{\rho} \left[ 1 - \frac{RT}{Mv^2} \left( \sqrt{1 + \frac{Mv^2}{3RT}} - 1 \right) \right]$$

where M= Molecular Weight,  $\rho$  = density and  $R=8.3143 \text{ JK}^{-1}\text{mol}^{-1}$ .

Ref: *A. Varada Rajulu and P. Mabu Sab*, Bull. Mater. Sci., Vol. 18 (June 1995), No. 3, pp. 247-253.

### 13. Wada Constant (W)

$$1. W = \frac{M}{\rho^{6/\gamma}} V^{2/\gamma} = M^{1/\gamma} R^{6/\gamma}$$

where  $\rho$  = density, M= Molecular Weight and R= Gas Constant.

Ref: *R.P. Singh, G.V. Reddy, S. Majumdar and Y.P. Singh*, J. Pure Appl. Ultrason, Vol. 5(1983), pp. 52-54.

$$2. W = (M. \beta^{-1/\gamma})/\rho$$

where  $\rho$  = density,  $\beta$  = adiabatic compressibility and M= molecular Weight.

Ref: *P.A.K. Ekka, G.V. Reddy and R.P. Singh*, Acustica, Vol. 46, (1980), No. 3, pp-341-342.

### 14. Space Filling Factor (r)

$$r = v/U$$

where U = represents the sound velocity when the entire volume of space is filled with molecules.

Ref: *A. Varada Rajulu, K. Chowdji Rao and S. Venkata Naidu*, Acustica, Vol. 75 (1991), pp. 213-216.

### 15. Relative Association ( $R_A$ )

$$R_A = (\rho / \rho_0) (v_0/v)^{1/3}$$

where  $\rho_0$  = density of solvent,  $v_0$  = ultrasonic velocity of solvent

Ref: *Anwar Ali and Anil Kumar Nain*, *Acoustics Letters*, Vol. 19 (1996), No. 9, pp. 181-187.

### 16. Optical Refractive Index(n)

$$v = [10(s/h)]n^{1/x}$$

where  $s/h$  = ratio of surface tension to the coefficient of viscosity,

$X = 1/L_f$

where  $l$  is mean free path of the molecule and  $L_f$  is intermolecular free length

Ref: *S. Durai and P Ramadoss*, *Indian Journal of Pure and Applied Physics*, Vol. 42 (2004), pp. 334-337.

### 17. Solvation Number ( $S_n$ )

$$S_n = \frac{M}{M_o} \left( 1 - \frac{\beta}{\beta_o} \right) \left( \frac{100-x}{x} \right)$$

where  $M$  and  $M_o$  are molecular weight of Solvent and Solution respectively,  $\beta$  and  $\beta_o$  are adiabatic compressibility of Solvent and Solution respectively and  $x$  is the number of grams of salt in 100g of the solution.

Ref: *R Ezhil Pavai, P Vasantharani and A N Kannappan*, *Indian Journal of Pure and Applied Physics*, Vol.42, , pp.934-936.

### 18. Attenuation ( $\alpha/f^2$ )

$$\alpha/f^2 = 8\pi^2 \eta / 3\rho v^3$$

where  $\alpha$  is absorption coefficient,  $f$  is frequency,  $\eta$  is shear viscosity,  $\rho$  is density and  $v$  is speed of sound in dispersion.

Ref: *A.Varada Rajulu, G.Sreenivasulu and K.S.Raghuraman*, *Indian Journal of Chemical Technology*, Vol.1, Sept 1994, pp.302-304.